

**REPORTABLE DISEASES: UTAH
1900 - 1909**

	1900	1901	1902	1903	1904	5 YR TOT	1905	1906	1907	1908	1909	5 YR TOT	10 YR AVG
BRUCELLOSIS (UNDULANT FEVER)						0		409	913	780	669	2,771	693
DIPHTHERIA	NO DATA AVAILABLE												
ENCEPHALITIS (ACUTE INFECTIONS)	NO DATA AVAILABLE												
INFLUENZA	NO DATA AVAILABLE												
MALARIA	NO DATA AVAILABLE												
MENINGITIS (EPIDEMIC)	NO DATA AVAILABLE												
PERTUSSIS (WHOOPIING COUGH)						0		1,748	1,691	1,279	2,670	7,388	1,847
PNEUMONIA						0		734	1,014	1,323	1,058	4,129	1,032
POLIO	NO DATA AVAILABLE												
RABIES HUMAN	NO DATA AVAILABLE												
ROCKY MT SPOTTED FEVER	NO DATA AVAILABLE												
RUBEOLA (MEASLES)						0		111	4,855	10,049	689	15,704	3,926
SCARLET FEVER						0		660	773	1,388	2,030	4,851	1,213
SMALLPOX						0		893	269	652	1,846	3,660	915
STREPTOCOCCAL INFECTIONS	NO DATA AVAILABLE												
TUBERCULOSIS (ALL FORMS)						0		159	158	217	228	762	191
TULAREMIA	NO DATA AVAILABLE												
TYPHOID FEVER						0		871	881	777	1,506	4,035	1,009
Data Source: Utah Morbidity and Mortality Statistics (Residents Only) 1906-1940, 1908-1944 and 1918-1956													
Summary Tables published by the Utah Department of Health in the Utah Vital Statistics Annual Reports of 1940, 1944 and 1956.													
Most figures are taken from the 1944 Report. Encephalitis, malaria and streptococcal data come from the 1954-56 reports.													
Epidemic meningitis figures match later data that is labeled as meningococcal meningitis.													
The 1955-56 table appear to include only resident cases; however, the tables are not labeled.													
Residents do not include military cases and residents of Indian Reservations.													