


# Central Utah Local Health District Report

## Inside

Introduction.....	1
Risk Factors.....	2
Mortality.....	5
Pre-transport.....	6
Hospitalization.....	7
Sources.....	8
Appendix A .....	9



## Introduction

Cardiovascular disease is the leading cause of death in the United States and Utah. Cardiovascular disease includes those diseases that affect the heart and blood vessels including coronary heart disease, stroke, congestive heart failure, and hypertension. Several modifiable risk factors contribute to cardiovascular disease including high blood pressure, high blood cholesterol, tobacco use, excessive alcohol consumption, inadequate physical activity, poor nutrition, obesity, and diabetes.

This report examines the burden of cardiovascular disease in Utah and each of Utah's 12 local health districts. The data presented on cardiovascular disease come from several sources including the Utah Behavioral Risk Factor Surveillance System, Utah Death Certificates, and the Utah Hospital Discharge Database.

The data have been divided by geographic area to provide a general overview of health status in each local health district. Local programs will find this information useful when presenting a case for heart disease and stroke prevention in their areas. The information is also useful for program planning, setting priorities, and evaluation.



Heart Disease & Stroke  
Prevention Program

**Table 1.****Demographics, Utah vs Central Utah Health District**

Demographic Subgroup	Health District Population		95% Confidence Intervals		State Population	
	Number of Persons	Distribution	Lower	Upper	Number of Persons	Distribution
<b>Gender</b>						
Males	36,108	50.6%			1,303,786	50.5%
Females	35,207	49.4%			1,278,585	49.5%
Total	71,315	100.0%			2,582,371	100.0%
<b>Age Group</b>						
0 to 4	5,724	8.0%			254,065	9.8%
5 to 17	15,843	22.2%			550,504	21.3%
18 to 49	31,435	44.1%			1,221,336	47.3%
50 or Over	18,313	25.7%			556,466	21.5%
Total	71,315	100.0%			2,582,371	100.0%
<b>Race/Ethnicity</b>						
White	66,334	97.1%			2,241,072	93.8%
Black	205	0.3%			22,534	0.9%
American Indian/Alaskan Native	899	1.3%			32,191	1.3%
Asian	336	0.5%			44,608	1.9%
Pacific Islander	210	0.3%			17,368	0.7%
Other	321	0.5%			31,266	1.3%
Total	68,305	100.0%			2,389,039	100.0%
Non-Hispanic	64,447	94.4%			2,135,966	89.4%
Hispanic	3,858	5.6%			253,073	10.6%
<b>Unemployment</b>	1,618	5.6%			56,610	4.7%
<b>Income</b>						
Less Than \$20,000	13,900	19.5%	16.3%	23.0%	390,900	15.1%
\$20,000-\$49,999	36,600	51.3%	47.0%	55.6%	1,054,600	40.8%
\$50,000 and Over	20,800	29.2%	25.6%	33.1%	1,136,800	44.0%
<b>Education</b>						
Less Than High School	5,800	8.1%	6.1%	10.8%	183,800	7.1%
H.S. Grad or G.E.D.	25,200	35.3%	31.6%	39.3%	717,500	27.8%
Some Post High School	26,400	37.0%	33.2%	41.1%	870,300	33.7%
College Graduate	13,900	19.5%	16.7%	22.6%	810,700	31.4%
<b>Do Not Have Health Care Coverage</b>	11,800	16.5%	13.6%	19.9%	364,900	14.1%
<b>Unable to Get Health Care Due to Cost</b>	10,100	14.2%	11.6%	17.2%	320,200	12.4%
<b>Report Having "Fair" or "Poor" Health</b>	11,600	16.2%	13.6%	19.3%	347,300	13.5%
<b>Seven or More Days of Poor Mental Health</b>	12,000	16.8%	14.0%	20.1%	392,800	15.2%

Sources: Population data from 2006 Governor's Office of Planning and Budget Population Estimates; race/ethnicity data from 2004 U.S. Census Bureau county estimates; unemployment data from 2004 Department of Workforce Services Unemployment Rates; income, education, health status, and health care coverage data from 2003-2005 BRFSS age-adjusted to 2000 population

# Risk Factors

## Physical Activity

Regular physical activity reduces the risk of cardiovascular disease in general and of coronary heart disease in particular. Adults should get at least 30 minutes of moderate intensity exercise five or more days a week, or at least 20 minutes of vigorous exercise three or more days a week.<sup>1</sup>

## Fruits and Vegetables

A diet of five or more servings of fruits and vegetables per day is associated with a reduced risk of coronary heart disease, some types of cancer, diabetes, and other chronic diseases.<sup>2</sup> Fruits and vegetables are low in fat and high in vitamins, minerals and fiber. A healthy diet that includes fruits and vegetables can reduce blood pressure and blood cholesterol levels. In general, a person should eat at least two servings of fruit and three servings of vegetables a day.

## Obesity

Obesity is fast approaching smoking as the leading preventable cause of death in the United States. An estimated 300,000 deaths a year can be attributed to obesity. Adults who are overweight or obese are at increased risk of hypertension, high blood cholesterol, type 2 diabetes, coronary heart disease, stroke, and certain forms of cancer.<sup>3,9</sup>

## Smoking


Smoking is the leading preventable cause of death in the United States. Cigarette smoking causes heart disease, chronic lung disease, several kinds of cancer, and other diseases.<sup>4</sup>

## Diabetes

Diabetes is a disease in which blood glucose levels are above normal. Diabetes can cause serious health complications including heart disease, blindness, kidney failure, and lower-extremity amputations. Diabetes is the sixth leading cause of death in the United States and Utah.

**Figure 1.**

Risk Factors. Percentage of Adults Aged 18 and Over, by Risk Factor, Utah vs Central Utah Health District



Source: BRFSS, Office of Public Health Assessment, Utah Department of Health 2002-2005, Age-adjusted to 2000 population

\* Includes 2001 data

\*\* Among adults who smoke


## High Blood Pressure

High blood pressure is a leading cause for cardiovascular disease. Hypertension, often called high blood pressure, is having a systolic blood pressure of 140mm Hg and higher or a diastolic blood pressure of 90mm Hg and higher. A person with high blood pressure should be diagnosed by a doctor before concluding that that person has hypertension. Hypertension is related to an increased incidence of heart disease, heart attack, stroke, atherosclerosis, and coronary artery disease. Individuals with high blood pressure should take steps to reduce and control their blood pressure.<sup>7</sup>

## High Blood Cholesterol

High levels of cholesterol and triglycerides increase the risk for heart disease. The National Heart, Lung, and Blood institute defines high blood cholesterol as 240 mg/dl or greater and "borderline high" cholesterol as 200 to 239 mg/dl. Cholesterol and other fats can't dissolve in the blood. They have to be transported to and from the cells by special carriers called lipoproteins. There are two types of lipoproteins that are important in our bodies: low-density lipoprotein (LDL), which carries cholesterol to the arteries, and high-density lipoprotein (HDL), which removes cholesterol from the arteries. High levels of LDL and low levels of HDL are related to an increased incidence of heart disease, including heart attack, atherosclerosis, and coronary artery disease. Individuals with high cholesterol should take steps to reduce and control their cholesterol levels.<sup>8</sup>


**Figure 2.**  
Doctor-diagnosed High Blood Pressure.  
Percentage of Adults Aged 18 and Over,  
Utah vs Central Utah Health District



Source: BRFSS, Office of Public Health Assessment, Utah Department of Health 2003-2005, Age-adjusted to 2000 population

\* Respondents who were told they had high blood pressure during pregnancy were excluded from the population

**Figure 3.**  
Doctor-diagnosed High Cholesterol.  
Percentage of Adults Aged 18 and Over,  
Utah vs Central Utah Health District




Source: BRFSS, Office of Public Health Assessment, Utah Department of Health 2003-2005, Age-adjusted to 2000 population

\* Percentage of all respondents, results should not be interpreted as prevalence of high cholesterol.

**Figure 4.**

Mortality. Rate per 100,000 Population, by Cause of Death,  
Utah vs Central Utah Health District



Source: Utah Death Certificate Database, Office of Vital Records, Utah Department of Health 2002-2004, Age-adjusted to 2000 population  
ICD-10 Codes : See appendix A


# Pre-transport

Early access to medical services is critical in determining the outcome of a stroke or heart attack. Only a small percent of patients who could benefit from treatments like clot-dissolving therapy or

defibrillation actually receive treatment early enough for them to work.<sup>5</sup> Individuals who do not recognize the signs and symptoms of a heart attack or stroke are less likely to receive the treatment they need before it's too late. The best strategy is to be aware of the early warning signs and respond to them by calling 911 immediately.

**Figure 5.**


Pre-transport Death and Knowledge of Signs and Symptoms of Stroke.  
Utah vs Central Utah Health District



Sources: \* Utah Death Certificate Database, Office of Vital Records, Utah Department of Health 1999-2004, Age-adjusted to 2000 population; defined as the percentage of stroke deaths that occur before transport to an emergency department  
\*\* BRFSS, Office of Public Health Assessment, Utah Department of Health 2001-2005, Age-adjusted to 2000 population; defined as the percentage of adults 18 and over who could identify all 5 of the major symptoms of stroke

**Figure 6.**

Pre-transport Death and Knowledge of Signs and Symptoms of Heart Attack.  
Utah vs Central Utah Health District



Sources: \* Utah Death Certificate Database, Office of Vital Records, Utah Department of Health 1999-2004, Age-adjusted to 2000 population; defined as the percentage of heart attack deaths that occur before transport to an emergency department  
\*\* BRFSS, Office of Public Health Assessment, Utah Department of Health 2001-2005, Age-adjusted to 2000 population; defined as the percentage of adults 18 and over who could identify all 5 of the major symptoms of heart attack

# Hospitalization

In 2004 the United States spent over \$225 billion on cardiovascular disease. Almost half of those costs were spent on hospitalization.<sup>6</sup>

**Table 2.**

**Average and Total Hospitalization Costs for Primary Diagnosis, Utah vs Central Utah Health District**

Condition	Central Utah Health District			Utah		
	Average Hospitalization Cost per Diagnosis	Average Total Annual Cost	Percent of Cost Paid by Government	Average Hospitalization Cost per Diagnosis	Average Total Annual Cost	Percent of Cost Paid by Government
All Cardiovascular Disease	\$18,499.98	\$15,502,979.67	72.4%	\$22,545.81	\$445,738,091.00	67.6%
Coronary Heart Disease	\$27,175.82	\$8,415,444.33	68.6%	\$28,346.39	\$212,777,462.67	63.4%
Acute Myocardial Infarction	\$31,009.74	\$3,473,091.33	67.7%	\$30,802.51	\$84,214,063.33	61.4%
Congestive Heart Failure	\$9,868.35	\$1,648,014.67	84.6%	\$15,843.41	\$46,616,597.33	79.7%
Stroke	\$10,638.35	\$1,195,041.33	72.2%	\$15,214.63	\$49,442,462.00	69.5%

Source: Utah Inpatient Hospital Discharge Data, Office of Health Care Statistics, Utah Department of Health, 2002-2004; Government costs include Medicaid, Medicare, CHIP, Worker's Compensation, and Other Government

**Table 3.**

**Average and Total Hospitalization Costs for Any Diagnosis,\* Utah vs Central Utah Health District**

Condition	Central Utah Health District			Utah		
	Average Hospitalization Cost per Diagnosis	Average Total Annual Cost	Percent of Cost Paid by Government	Average Hospitalization Cost per Diagnosis	Average Total Annual Cost	Percent of Cost Paid by Government
All Cardiovascular Disease	\$14,990.12	\$40,603,236.67	74.5%	\$18,264.67	\$1,201,212,464.67	68.3%
Coronary Heart Disease	\$18,464.72	\$14,691,759.67	75.7%	\$20,710.43	\$407,463,960.00	71.4%
Acute Myocardial Infarction	\$31,923.67	\$4,426,749.00	70.1%	\$31,635.96	\$115,439,636.00	65.1%
Congestive Heart Failure	\$15,376.61	\$12,752,334.00	85.9%	\$20,690.93	\$306,446,464.00	81.0%
Stroke	\$13,463.44	\$2,912,590.67	72.7%	\$17,720.93	\$106,502,764.33	70.6%

Source: Utah Inpatient Hospital Discharge Data, Office of Health Care Statistics, Utah Department of Health, 2002-2004; Government costs include Medicaid, Medicare, CHIP, Worker's Compensation, and Other Government

\* Defined as mention in either primary or secondary diagnoses codes

**Table 4.**

**Per Capita Costs for Cardiovascular Disease, Utah vs Central Utah Health District**


	Central Utah Health District	Utah
Amount Spent Per Capita on Prevention	\$0.99	\$0.21
Hospitalization Costs Per Capita (Primary Diagnosis)	\$222.43	\$184.67
Hospitalization Costs Per Capita (Any Diagnosis)*	\$582.56	\$497.66

Sources: Prevention costs calculated using local health department contract summary data Fiscal Year 2006, Heart Disease and Stroke Prevention Program; Hospitalization costs calculated from Utah Inpatient Hospital Discharge Data, Office of Health Care Statistics, Utah Department of Health, 2002-2004

\* Defined as mention in either primary or secondary diagnoses codes

**Figure 7.**

Hospitalization Rates. Rate per 10,000 Population, by Primary Diagnosis, Utah vs Central Utah Health District



Source: Utah Inpatient Hospital Discharge Data, Office of Health Care Statistics, Utah Department of Health, 2002-2004, Age-adjusted to 2000 population  
ICD-9 Codes : See appendix A

## Sources

1. U.S. Department of Health and Human Services. Physical Activity and Health: A Report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion 1996.
2. U.S. Department of Health and Human Services and U.S. Department of Agriculture. Dietary Guidelines for Americans, 2005. 6th Edition, Washington, DC: U.S. Government Printing Office, January 2005.
3. U.S. Department of Health and Human Services. The Surgeon General's call to action to prevent and decrease overweight and obesity. [Rockville, MD]: U.S. Department of Health and Human Services, Public Health Service, Office of the Surgeon General; [2001]. Available from: U.S. GPO, Washington.
4. U.S. Department of Health and Human Services. The Health Benefits of Smoking Cessation. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. DHHS Publication No. (CDC) 90-8416. 1990.
5. U.S. Department of Health and Human Services. Healthy People 2010: Understanding and Improving Health. 2nd ed. Washington, DC: U.S. Government Printing Office, November 2000.
6. National Heart, Lung, and Blood Institute. Morbidity & mortality: 2004 chart book on cardiovascular, lung, and blood diseases. U.S. Department of Health and Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute; 2004.
7. Sheridan S, Pignone M, Donahue K. Screening for High Blood Pressure: A Review of the Evidence for the U.S. Preventive Services Task Force. *Am J Prev Med.* 2003;25:151-158.
8. Pignone MP, Phillips CJ, Atkins D, Teutsch SM, Mulrow CD, Lohr KN. Screening and treating adults for lipid disorders. *Am J Prev Med.* 2001;20:77-89.
9. Allison DB, Fontaine KR, Manson JE, Stevens J, VanItallie TB. Annual deaths attributable to obesity in the United States. *JAMA* 1999 Oct 27;282(16):1530-8.

# Appendix A

The International Classification of Diseases, Ninth and Tenth Revisions, Clinical Modification (ICD-

9-CM and ICD-10 CM) were used to identify the specific disease categories in this report. The following ICD-9 or ICD-10 codes have been grouped together to determine rates for the specific conditions included in this report.

ICD-9 and ICD-10 Codes by Cause

Condition/Procedure	Type of Data	ICD-9 Codes	ICD-10 Codes
All Cardiovascular Disease	Mortality	390-448	I00-I78, G45
	Hospital Discharge	390-448	
Coronary Heart Disease	Mortality	402, 410-414, 429.2	I11, I20-I25
	Hospital Discharge	402, 410-414, 429.2	
Acute Myocardial Infarction	Mortality	410	I21-I22
	Hospital Discharge	410	
Congestive Heart Failure*	Mortality	428.0, 428.1, 428.9	I50.0, I50.1, I50.9
	Hospital Discharge	428.0, 428.1, 428.9, 398.91**	
Stroke	Mortality	430-438	I60-I69
	Hospital Discharge	430-438	

\* According to guidelines governing the writing of cause-of-death statements, congestive heart failure should not be listed as an underlying cause of death and, therefore, these data should be interpreted with caution.  
 \*\* Mortality data were not coded to the fifth digit and, therefore, ICD-9 code 398.91 was not included in the definition of congestive heart failure when analyzing mortality data.

# Notes

A large rectangular area with horizontal dotted lines for writing notes.

# Notes

A large rectangular area with a solid top border and a solid right border. The interior is filled with horizontal dotted lines, providing a space for writing notes.