

R430-60-16: INFECTION CONTROL.

- (1) Staff shall wash their hands thoroughly with liquid soap and warm running water at the following times:
- (a) before handling or preparing food or bottles;
 - (b) before and after eating meals and snacks or feeding children;
 - (c) before and after diapering a child;
 - (d) after using the toilet or helping a child use the toilet;
 - (e) before administering medication;
 - (f) after coming into contact with body fluids;
 - (g) after playing with or handling animals;
 - (h) when coming in from outdoors; and
 - (i) after cleaning or taking out garbage.

Rationale / Explanation

Handwashing is the most important way to reduce the spread of infection. Many studies have shown that unwashed or improperly washed hands are the primary carriers of infection. Deficiencies in handwashing have contributed to many outbreaks of diarrhea among children and caregivers in child care centers. In centers that have implemented a handwashing training program, the incidents of diarrheal illness has decreased by 50%. One study also found that handwashing helped to reduce colds when frequent proper handwashing practices were incorporated into a child care center's curriculum. CFOC, 3rd Ed. pgs. 110-111 Standard 3.2.2.1, pg. 114 Standard 3.2.3.1

Washing hands after eating is especially important for children who eat with their hands, to decrease the amount of saliva (which may contain organisms) on their hands. Good handwashing after playing in sandboxes will help prevent ingesting parasites that can be present in contaminated sand and soil. Animals, including pets, are a source of infection for people and people may be a source of infection for animals. CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2.1.

Illness can be spread in a variety of ways that can be reduced with proper handwashing, including:

- *in human waste (urine, stool)*
 - *in body fluids (saliva, nasal discharge, secretions from open injuries, eye, discharge, blood, etc.)*
 - *through cuts or skin sores*
 - *by direct skin-to-skin contact*
 - *by touching an object that has germs on it*
 - *in drops of water that travel through the air, such as those produced by sneezing or coughing.*
- CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2.1*

Since many infected people carry communicable diseases without having symptoms and many are contagious before they experience a symptom, staff members need to protect both themselves and children by following good hygiene practices on a routine basis. CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2

Running water over the hands removes soil, including infection-causing organisms. Wetting the hands before applying soap helps create a lather. The soap lather loosens soil and brings it into the solution on the surface of the skin. Rinsing the lather off into a sink removes the soil from the hands that the soap loosened. Warm water (no less than 60 degrees Fahrenheit and no more than 120 degrees) is more comfortable than cold water, which increases the likelihood that children and adults will adequately rinse their hands.

Using liquid soap is preferable over bar soap. Bar soaps sitting in water have been shown to be heavily contaminated with pseudomonas and other bacteria. In addition, many children do not have the dexterity to

R430-60-16: INFECTION CONTROL.

handle a bar of soap, and many adults and children do not take the time to rise off the soil that has gotten on the bar of soap before putting it down. CFOC, 3rd Ed. pgs. 258-259 Standard 5.6.0.3

Using a paper towel to turn off the faucet after handwashing can prevent the re-contamination of just-washed hands by germs on the faucet. CFOC, 3rd Ed. pgs. 111-112 Standard 3.2.2.2

Enforcement

Water will be considered warm if it is between 60 degrees and 120 degrees Fahrenheit.

Hand sanitizers may be used to meet the hand washing requirements for adults and children age 2 and older without visibly dirty hands.

Level 2 Noncompliance if handwashing does not take place after a caregiver uses the toilet.

Level 3 Noncompliance otherwise.

- (2) **The provider shall ensure that children wash their hands thoroughly with liquid soap and warm running water at the following times:**
- (a) **before and after eating meals and snacks;**
 - (b) **after using the toilet;**
 - (c) **after coming into contact with body fluids;**
 - (d) **after playing with animals; and**
 - (e) **when coming in from outdoors.**

Rationale / Explanation

Handwashing is the most important way to reduce the spread of infection. Many studies have shown that unwashed or improperly washed hands are the primary carriers of infection. Deficiencies in handwashing have contributed to many outbreaks of diarrhea among children and caregivers in child care centers. In centers that have implemented a handwashing training program, the incidents of diarrheal illness has decreased by 50%. One study also found that handwashing helped to reduce colds when frequent proper handwashing practices were incorporated into a child care center's curriculum. CFOC, 3rd Ed. pgs. 110-111 Standard 3.2.2.1, pg. 114 Standard 3.2.3.1

Washing hands after eating is especially important for children who eat with their hands, to decrease the amount of saliva (which may contain organisms) on their hands. Good handwashing after playing in sandboxes will help prevent ingesting parasites that can be present in contaminated sand and soil. Animals, including pets, are a source of infection for people and people may be a source of infection for animals. CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2.1

Illness can be spread in a variety of ways that can be reduced with proper handwashing, including:

- in human waste (urine, stool)
- in body fluids (saliva, nasal discharge, secretions from open injuries, eye, discharge, blood, etc.)
- through cuts or skin sores

R430-60-16: INFECTION CONTROL.

- *by direct skin-to-skin contact*
- *by touching an object that has germs on it*
- *in drops of water that travel through the air, such as those produced by sneezing or coughing.*
CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2.1 .

Since many infected people carry communicable diseases without having symptoms and many are contagious before they experience a symptom, staff members need to protect both themselves and children by following good hygiene practices on a routine basis. CFOC, 3rd Ed. pgs. 100-111 Standard 3.2.2.1

Running water over the hands removes soil, including infection-causing organisms. Wetting the hands before applying soap helps create a lather. The soap lather loosens soil and brings it into the solution on the surface of the skin. Rinsing the lather off into a sink removes the soil from the hands that the soap loosened. Warm water (no less than 60 degrees Fahrenheit and no more than 120 degrees) is more comfortable than cold water, which increases the likelihood that children and adults will adequately rinse their hands.

*Using liquid soap is preferable over bar soap. Bar soaps sitting in water have been shown to be heavily contaminated with *Pseudomonas* and other bacteria. In addition, many children do not have the dexterity to handle a bar of soap, and many adults and children do not take the time to rise off the soil that has gotten on the bar of soap before putting it down. CFOC, 3rd Ed. pgs. 258-259 Standard 5.6.0.3*

Using a paper towel to turn off the faucet after handwashing can prevent the re-contamination of just-washed hands by germs on the faucet. 3rd Ed. pgs. 111-112 Standard 3.2.2.2

Enforcement

Water will be considered warm if it is between 60 degrees and 120 degrees Fahrenheit.

Hand sanitizers may be used to meet the hand washing requirements for adults and children age 2 and older without visibly dirty hands.

Always Level 2 Noncompliance.

- (3) Only single use towels from a covered dispenser or an electric hand-drying device may be used to dry hands.**

Rationale / Explanation

Shared hand drying towels can transmit infectious disease. Preventing shared use of individual towels assigned to a single child is difficult. The use of a cloth towel roller is not recommended for two reasons. First, children often use cloth roll dispensers improperly, resulting in more than one child using the same section of towel. And second, incidents of accidental strangulation in these devices have been reported. CFOC, 3rd Ed. pgs. 111-112 Standard 3.2.2.2

Enforcement

R430-60-16: INFECTION CONTROL.

This rule does not apply to paper towels used for activities other than drying hands, such as cleaning up spills.

Always Level 3 Noncompliance.

- (4) The provider shall ensure that toilet paper is accessible to children, and that it is kept on a dispenser.**

Rationale / Explanation

The purpose of this rule to prevent the spread of disease through fecal matter. If toilet paper is not on a dispenser, children pick it up with hands that may be contaminated with fecal matter, which remains on the roll and is transferred to the next child when he or she picks the roll up. CFOC, 3rd Ed. pgs. 258-259 Standard 5.6.0.3

Enforcement

Toilet paper is not considered accessible unless the child can reach it while he/she is sitting on the toilet.

Level 2 Noncompliance if a toilet has no toilet paper and there are no spare rolls of toilet paper available in the facility or if toilet paper is not kept on a dispenser.

Level 3 Noncompliance If a toilet has no toilet paper but there are spare rolls of toilet paper available in the facility.

- (5) The provider shall clean and sanitize all washable toys and materials weekly, or more often if necessary.**

Rationale / Explanation

Contamination of toys and other objects in child care areas plays a role in the transmission of disease in child care settings. All toys can spread disease when children touch the toys after putting their hands in their mouth during play or eating, or after toileting with inadequate handwashing. Using a mechanical dishwasher is an acceptable labor-saving approach for plastic toys as long as the dishwasher can wash and sanitize the surfaces. CFOC, 3rd Ed. pgs. 116-118, Standards 3.3.0.1, 3.3.0.2

Enforcement

This rule is out of compliance if toys or materials are visibly dirty during an inspection or if providers indicate they do not clean and sanitize all washable toys and materials at least weekly. Since toys in child care settings are heavily used, every toy is not expected to be perfectly clean all the time.

Always Level 3 Noncompliance.

- (6) Stuffed animals, cloth dolls, and dress-up clothes must be machine washable. Pillows must be machine washable, or have removable covers that are machine washable. The provider shall wash stuffed animals, cloth dolls, dress-up clothes, and pillows or covers weekly.**

Rationale / Explanation

Contamination of toys and other objects in child care areas plays a role in the transmission of disease in child

R430-60-16: INFECTION CONTROL.

care settings. All toys can spread disease when children touch the toys after putting their hands in their mouth during play or eating, or after toileting with inadequate handwashing. CFOC, 3rd Ed. pgs. 116-118, Standards 3.3.0.1, 3.3.0.2

Many allergic children have allergies to dust mites, which are microscopic insects that ingest the tiny particles of skin that people shed normally every day. Dust mites live in fabric, but can be killed by frequent washing and drying in a heated dryer. CFOC, 3rd Ed. pg. 239 Standard 5.3.1.4

Lice, scabies, and ringworm can also be spread through fabrics. CFOC, 3rd Ed. pg. 118 Standard 3.3.0.4, pg. 136 Standard 3.6.1.5

Enforcement

This rule is out of compliance if these items are visibly dirty during an inspection or if providers indicate that they do not wash them at least weekly.

Large stuffed animals meant to be used as pillows need to be machine washable or have removable covers that are machine washable.

Level 2 Noncompliance if an item is visibly dirty with feces.

Level 3 Noncompliance otherwise.

(7) If water play tables or tubs are used, they shall be washed and sanitized daily, and children shall wash their hands prior to engaging in the activity.

Rationale / Explanation

The purpose of this rule is to avoid the spread of disease as multiple children's hands play in the water in water tables. Contamination of hands, toys, and equipment in the room where water play tables are located plays a role in the transmission of disease in child care settings. CFOC, 3rd Ed. pg. 275 Standard 6.2.4.2.

Enforcement

Hand sanitizers may be used to meet the hand washing requirements for adults and children age 2 and older without visibly dirty hands.

Always Level 3 Noncompliance.

(8) Persons with contagious TB shall not work or volunteer in the center.

Rationale / Explanation

The purpose of this rule is to prevent the spread of TB from infected adults to children. Young children acquire TB from infected adults or adolescents. Tuberculosis organisms are spread by the inhalation of small particles which are produced when an infected adult or adolescent coughs or sneezes. Transmission usually occurs in an indoor environment. CFOC, 3rd Ed. pgs. 39-40 Standard 1.7.0.1

R430-60-16: INFECTION CONTROL.

Enforcement

Always Level 1 Noncompliance.

- (9) **Children's clothing which is wet or soiled from body fluids:**
- (a) shall not be rinsed or washed at the center; and
 - (b) shall be placed in a leakproof container, labeled with the child's name, and returned to the parent.

Rationale / Explanation

Containing and minimizing the handling of soiled clothing so it does not contaminate other surfaces is essential to prevent the spread of infectious disease. Rinsing soiled clothing or putting stool into a toilet in the child care center increases the likelihood that other surfaces will be contaminated. CFOC, 3rd Ed. pgs. 108-110 Standard 3.2.1.5.

Enforcement

Plastic grocery bags may be used for wet or soiled clothing if they don't have holes in the bottom or sides. Grocery bags with holes in the bottom or sides cannot be used, because they are not leakproof. A bag without holes that still leaks when holding wet or soiled clothes cannot be used.

A provider may meet the intent of this rule by putting soiled clothing in a leakproof container and then putting the container in that child's diaper bag or cubbie.

When a child has a bathroom accident and the sheets and/or the clothing used belong to the Licensee, the center staff can and should wash the sheets and/or the clothing. The rule prohibiting washing the clothing only pertains to clothing belonging to the children. Staff can also take soiled clothing home for laundering.

Level 2 Noncompliance if children's clothing soiled with fecal matter is rinsed or washed at the center.

Level 3 Noncompliance otherwise.

- (10) **If the center uses a potty chair, the provider shall clean and sanitize the chair after each use.**

Rationale / Explanation

The purpose of this rule is to prevent the spread of disease through fecal matter or the growth of disease-causing microorganisms in urine or stool that sit in potty chairs over time. It is also necessary in order to prevent naturally curious toddlers from playing in urine or feces that may be in potty chairs after they are used. CFOC, pg. 105 Standard 3.029.

Because of the difficulties in the sanitary handling of potty chairs, the American Academy of Pediatrics and the American Public Health Association recommend that they not be used.

Enforcement

R430-60-16: INFECTION CONTROL.

A toilet training seat is only considered a potty chair if it collects and holds urine or feces. Toddler toilet seats that are placed over the regular toilet seat, where the urine and feces are flushed, are not considered to be potty chairs.

Always Level 2 Noncompliance.

- (11) The center shall have a portable body fluid clean up kit.**
- (a) All staff shall know the location of the kit and how to use it.**
 - (b) The provider shall use the kit to clean up spills of body fluids.**
 - (c) The provider shall restock the kit as needed.**

Rationale / Explanation

Children and adults may unknowingly be infected with infectious agents such as hepatitis B, HIV, or other infectious agents found in blood. Blood and body fluids containing blood (such as water discharges from injuries) pose the highest potential risk because bloody body fluids contain the highest concentration of viruses. In addition, the hepatitis B virus can survive in a dried state for at least a week and perhaps even longer. Some other body fluids such as saliva contaminated with blood or blood-associated fluids may contain live viruses but at lower concentrations than are found in blood itself. Many other types of infectious germs may be contained in human waste and other body fluids. Because many people carry such communicable diseases without having symptoms, and many are contagious before they experience symptoms, adults and children alike need to be protected by following safe procedures for handling body fluids. CFOC, 3rd Ed. pgs. 30-31 Standard 1.4.5.3 pgs. 114-116 Standard 3.2.3.4

Suggested contents for a body fluid clean up kit include:

- (1) disposable gloves;*
- (2) clumping cat litter, sawdust, or other absorbent material;*
- (3) plastic garbage bags with ties or fasteners;*
- (4) a plastic scoop and dustpan, or other tools to clean up absorbed body fluids;*
- (5) paper towels; and*
- (6) disinfectant.*

See CFOC, 3rd Ed. pg. 444 Appendix for an instruction page on proper clean up of body fluids. See CFOC, 3rd Ed. pg. 428 Appendix D for information on removing disposable gloves after cleaning up body fluids.

Enforcement

Droplets of body fluid are not considered a "spill" of body fluids.

Level 2 Noncompliance if a body fluid spill is not properly cleaned up or no one at the center knows the location of the kit or how to properly use it.

Level 3 Noncompliance otherwise.

- (12) The provider shall notify the local health department, on the day of discovery, of any reportable**

R430-60-16: INFECTION CONTROL.

infectious diseases among children or caregivers, or any sudden or extraordinary occurrence of a serious or unusual illness, as required by the local health department.

Rationale / Explanation

Reporting infectious disease to the local health department provides the department with knowledge of illnesses within the community and allows them to offer preventive measures to children and families exposed to an outbreak of disease. CFOC, 3rd Ed. pg. 355 Standard 9.2.3.3

The following is a sample of diseases which may be required to be reported to local health departments. Providers should check with the local health department in their area for exact reporting requirements.

- Chickenpox
- Diarrheal diseases, if two or more children or staff members in one classroom experience diarrhea within a 48 hour period.
- Diphtheria
- Giardiasis
- Hepatitis A, B, and C
- HIV and AIDS
- Influenza
- Measles
- Meningococcal infections
- Mumps
- Rubella
- Sexually transmitted diseases
- Shigellosis
- Viral Meningitis
- Whooping Cough

A free informational guide for controlling communicable diseases in child care centers can be found and printed at: <http://health.utah.gov/epi/cdepi/daycarebook.pdf>

This guide is published by the Utah Department of Health, Office of Epidemiology. It includes:

1. A parent letter that can be copied and filled in whenever a provider needs to notify parents of a communicable disease.
2. A one page fact sheet on each disease that can be posted at the center, given to parents, or used in staff training. These fact sheets include
 1. Incubation period
 2. Signs and symptoms
 3. Methods of transmission
 4. Minimum control measures
 5. Guidelines for exclusion of ill children from a child care program
 6. Information on body substance clean up
 7. A listing of all local health departments.

R430-60-16: INFECTION CONTROL.

Enforcement

Always Level 3 Noncompliance.

- (13) The provider shall post a parent notice at the center when any staff or child has an infectious disease or parasite.
- (a) The provider shall post the notice in a conspicuous location where it can be seen by all parents.
 - (b) The provider shall post and date the notice the same day the disease or parasite is discovered, and the notice shall remain posted for at least 5 days.

Rationale / Explanation

Notification of parents also allows them to closely observe their child for early signs and symptoms of illness. Early identification and treatment of infectious disease are important in reducing further transmission of the disease. CFOC, 3rd Ed. pg. 145 Standard 3.6.4.2

The purpose for leaving the notice posted for 5 days is so that parents of children who do not attend every day see the notice.

When posting notices, the center should be careful to maintain confidentiality by not posting the names of sick children. The informational guide found at: <http://health.utah.gov/epi/cdepi/daycarebook.pdf> contains a parent letter that can be copied and filled in whenever a provider needs to notify parents of a communicable disease, and a one page fact sheet on each disease that can be posted at the center and given to parents.

Enforcement

Posting the notice of illness on a computerized sign-in program so that all parents automatically see it when they sign their children in and out meets the requirement of this rule.

Always Level 2 Noncompliance.