

## Strategies for Tapering & Weaning

### Strategies for tapering:

From a medical standpoint, weaning from opioids can be done safely by slowly tapering the opioid dose and taking into account the following issues:

- A decrease by 10% of the original dose per week is usually well tolerated with minimal physiological adverse effects. Some patients can be tapered more rapidly without problems (over 6 to 8 weeks).
- If opioid abstinence syndrome is encountered, it is rarely medically serious although symptoms may be unpleasant.
- Symptoms of an abstinence syndrome, such as nausea, diarrhea, muscle pain and myoclonus can be managed with clonidine 0.1 – 0.2 mg orally every 6 hours or clonidine transdermal patch 0.1mg/24hrs (Catapres TTS-1™) weekly during the taper while monitoring for often significant hypotension and anticholinergic side effects. In some patients it may be necessary to slow the taper timeline to monthly, rather than weekly dosage adjustments.
- Symptoms of mild opioid withdrawal may persist for six months after opioids have been discontinued.
- Consider using adjuvant agents, such as antidepressants to manage irritability, sleep disturbance or antiepileptics for neuropathic pain.
- Do not treat withdrawal symptoms with opioids or benzodiazepines after discontinuing opioids.
- Referral for counseling or other support during this period is recommended if there are significant behavioral issues.
- Referral to a pain specialist or chemical dependency center should be made for complicated withdrawal symptoms.

### Recognizing and managing behavioral issues during opioid weaning:

Opioid tapers can be done safely and do not pose significant health risks to the patient. In contrast, extremely challenging behavioral issues may emerge during an opioid taper.

Behavioral challenges frequently arise in the setting of a prescriber who is tapering the opioid dose and a patient who places great value on the opioid he/she is receiving. In this setting, some patients will use a wide range of interpersonal strategies to derail the opioid taper. These may include:

- Guilt provocation (“You are indifferent to my suffering”)
- Threats of various kinds
- Exaggeration of their actual suffering in order to disrupt the progress of a scheduled taper

There are no fool-proof methods for preventing behavioral issues during an opioid taper, but strategies implemented at the beginning of the opioid therapy are most likely to prevent later behavioral problems if an opioid taper becomes necessary.